

Causal dissipative hydrodynamics obtained from the nonextensive/dissipative correspondence

Takeshi Osada^{1,*} and Grzegorz Wilk^{2,†}

¹*Theoretical Physics Laboratory, Faculty of Knowledge Engineering,
Musashi Institute of Technology, Setagaya-ku, Tokyo 158-8557, Japan*

²*The Andrzej Soltan Institute for Nuclear Studies, Hoza 69, 00681, Warsaw, Poland*
(Dated: October 29, 2008)

We derive the constitutive equations of causal relativistic dissipative hydrodynamics (d -hydrodynamics) from perfect nonextensive hydrodynamics (q -hydrodynamics) using the nonextensive/dissipative correspondence (NexDC) proposed by us recently. The q -hydrodynamics can be thus regarded as a possible model for the d -hydrodynamics facilitating its application to high energy multiparticle production processes. As an example we have shown that applying the NexDC to the perfect $1+1$ q -hydrodynamics, one obtains a proper time evolution of the bulk pressure and the Reynolds number.

PACS numbers: 24.10.Nz, 25.75.-q

The ideal hydrodynamic model successfully reproduces most of the RHIC experimental data [1] suggesting that matter created there resembles a strongly interacting quark-gluon fluid [2] rather than a free parton gas. However, there are also hints that, after all, a hadronic fluid cannot be totally ideal [3] and should be described by some kind of dissipative hydrodynamic model (or d -hydrodynamics) [4, 5, 6]. A number of such models have been proposed recently [7, 8, 9, 10, 11]. Although very promising, they have many difficulties, both in what concerns their proper formulation and applications [5, 10, 12]. Any new attempt to understand them more deeply is therefore welcome. In this letter we demonstrate that one can obtain constitutive equations of causal d -hydrodynamics starting from the relativistic perfect nonextensive hydrodynamics (q -hydrodynamics) supplemented by the nonextensive/dissipative correspondence (NexDC) proposed in [13] and taking place between the perfect q -hydrodynamics (based on the nonextensive Tsallis statistics [14] and described by the nonextensivity parameter q) and the usual d -hydrodynamics (based on the extensive Boltzmann-Gibbs statistics (BG) to which Tsallis statistics converges for $q \rightarrow 1$). We argue that the d -hydrodynamics emerging from such q -hydrodynamics is causal and preserves the simplicity of the latter which facilitates numerical applications [13] (instead of solving very complicated second order equations of d -hydrodynamics one can solve the much simpler equation of motion of the q -hydrodynamics).

Our q -hydrodynamics [13] is based on the relativistic nonextensive kinetic theory proposed in [15] (see also [16]). It is understood that such a theory accounts, by means of the parameter q , for all kinds of possible strong intrinsic fluctuations and long-range correlations existing in a hadronizing system. It replaces the usual notion of local thermal equilibrium by a kind of stationary state which also includes some interactions. It is described by the Tsallis nonextensive statistics [14] and characterized

by the nonextensive parameter $q > 1$ [17, 18, 19]. In this approach the *perfect q -hydrodynamic equations* (i.e., equations for the perfect nonextensive q -fluid without any additional currents) are given by the q -version of the energy momentum tensor:

$$\mathcal{T}_{q;\mu}^{\mu\nu} = [\varepsilon_q(T_q)u_q^\mu u_q^\nu - P_q(T_q)\Delta_q^{\mu\nu}]_{;\mu} = 0. \quad (1)$$

Here $\varepsilon_q(T_q)$, $P_q(T_q)$, $T_q = T_q(x)$ and $u_q^\mu(x)$ are, respectively, the nonextensive energy density, pressure and temperature field [13] and an accompanying hydrodynamic q -flow four vector field (which allows decomposition (1) to be performed; $\Delta_q^{\mu\nu} \equiv g^{\mu\nu} - u_q^\mu u_q^\nu$). If $T(x)$ and $u^\mu(x)$ are the corresponding temperature and velocity fields in the case of BG statistics ($q = 1$) with ε and P being the usual energy density and pressure then, as proposed in [13], one can map a q -flow into some dissipative d -flow by requiring that the following relations hold:

$$P(T) = P_q(T_q), \quad \varepsilon(T) = \varepsilon_q(T_q) + 3\Pi, \quad (2)$$

where Π is to be regarded as the bulk pressure of d -hydrodynamics which is defined by

$$\Pi \equiv \frac{1}{3}w_q[\gamma^2 + 2\gamma]. \quad (3)$$

Here $\delta u_q^\mu \equiv u_q^\mu - u^\mu$ is the four velocity difference between the non-extensive parameter $q > 1$ and $q = 1$ flows, $\gamma(x) \equiv \delta u_q^\mu(x)u_\mu(x) = -\frac{1}{2}\delta u_{q\mu}\delta u_q^\mu$ and $w_q \equiv \varepsilon_q + P_q$ is the q -enthalpy. If $T(x)$ and $\gamma(x)$ satisfy Eq. (2) relations in the whole space-time region, one can always transform equation of perfect q -hydrodynamics Eq. (1) into the following equation of d -hydrodynamics [4, 5, 6, 7, 8, 9, 10, 11]:

$$[\varepsilon(T)u^\mu u^\nu - [P(T) + \Pi]\Delta^{\mu\nu} + 2W^{(\mu}u^{\nu)} + \pi^{\mu\nu}]_{;\mu} = 0, \quad (4)$$

in which one recognizes the energy flow vector W^μ and the (symmetric and traceless) shear pressure tensor $\pi^{\mu\nu}$,

$$W^\mu = w_q[1 + \gamma]\Delta_\sigma^\mu \delta u_q^\sigma, \quad \pi^{\mu\nu} = w_q \delta u_q^{<\mu} \delta u_q^{\nu>} \quad (5)$$

(where $\Delta^{\mu\nu} \equiv g^{\mu\nu} - u^\mu u^\nu$, $A^{(\mu} B^{\nu)} \equiv \frac{1}{2}(A^\mu B^\nu + A^\nu B^\mu)$, and $a^{<\mu} b^{>} \equiv [\frac{1}{2}(\Delta_\lambda^\mu \Delta_\sigma^\nu + \Delta_\sigma^\mu \Delta_\lambda^\nu) - \frac{1}{3}\Delta^{\mu\nu} \Delta_{\lambda\sigma}] a^\lambda b^\sigma$). The d -hydrodynamics represented by Eq. (4) can be therefore regarded as a viscous counterpart of the perfect q -hydrodynamics represented by Eq. (1). We call this relation the *nonextensive/dissipative correspondence*, (NexDC). With the bulk pressure Eq. (3) and NexDC relations Eq. (2) one gets the q -enthalpy $w_q = w/[1 + \gamma]^2$ with $w \equiv Ts = \varepsilon + P$ being the usual enthalpy in BG statistics. Accordingly, the bulk pressure Eq. (3) can be written as

$$\Pi = w\Gamma, \quad \text{where} \quad \Gamma \equiv \frac{1}{3} \frac{\gamma(\gamma + 2)}{(\gamma + 1)^2}. \quad (6)$$

Finally, the NexDC leads to the following relations between components of the dissipative tensor:

$$W^\mu W_\mu = -3\Pi w, \quad \pi^{\mu\nu} W_\nu = -2\Pi W^\mu, \quad \pi_{\mu\nu} \pi^{\mu\nu} = 6\Pi^2. \quad (7)$$

We shall now address the central question of this Letter: is d -hydrodynamics (4) obtained in this way causal, as is the one discussed in [7, 8, 9, 10, 11]? To this end, let us first notice that, because we expect that $q - 1 \ll 1$ [13], our stationary state described by Eqs. (1) and (4) can be regarded as a *near equilibrium state* (with energy momentum tensor $\mathcal{T}_q^{\mu\nu} \equiv (\varepsilon_q + P_q)u_q^\mu u_q^\nu - P_q g^{\mu\nu}$). Accordingly, the state with $q = 1$, i.e., the one with no residual correlations between fluid elements and no intrinsic fluctuations present, is a *true equilibrium state* (with energy momentum tensor $\mathcal{T}_{\text{eq}}^{\mu\nu} \equiv \varepsilon(T)u^\mu u^\nu - P(T)\Delta^{\mu\nu}$ and with the equilibrium distribution given by the usual Boltzmann distribution function, $f_{\text{eq}}(x, p) = \exp[-p^\mu u_\mu(x)/k_B T(x)]$). Both states are near to each other and $\mathcal{T}_q^{\mu\nu} = \mathcal{T}_{\text{eq}}^{\mu\nu} + \delta\mathcal{T}^{\mu\nu}$, where $\delta\mathcal{T}^{\mu\nu} = -\Pi\Delta^{\mu\nu} + 2W^{(\mu} u^{\nu)} + \pi^{\mu\nu}$. As an immediate consequence of this, the conservation of q -entropy assumed in the ideal q -hydrodynamics, $[s_q u_q^\mu]_{;\mu} = 0$, results in the production of the usual entropy in d -hydrodynamics:

$$T[su^\mu]_{;\mu} = -\Pi u_{;\mu}^\mu - W_{;\mu}^\mu + W^\mu \frac{du_\mu}{d\tau} + \pi^{\mu\nu} u_{\langle\mu;\nu\rangle} \quad (8)$$

($\frac{dX}{d\tau} \equiv u^\mu X_{;\mu}$ is the proper time derivative acting on a quantity X). Acting by $d/d\tau$ on both sides of Eq. (6) and multiplying by Π , one obtains

$$-\Pi \left[\frac{d\Pi}{d\tau} - w \frac{d\Gamma}{d\tau} - \Gamma \frac{dw}{d\tau} \right] \equiv 0. \quad (9a)$$

Similarly, acting $d/d\tau$ on both sides of the first and the last terms in Eq. (7) one gets two other identities:

$$\frac{2}{3} \Gamma W_\mu \frac{dW^\mu}{d\tau} + \Pi \frac{d\Pi}{d\tau} + \Pi \Gamma \frac{dw}{d\tau} \equiv 0, \quad (9b)$$

$$-\frac{1}{6} \pi^{\mu\nu} \frac{d\pi_{\mu\nu}}{d\tau} + \Pi \frac{d\Pi}{d\tau} \equiv 0. \quad (9c)$$

Finally, acting by ∇_μ (where $\nabla_\mu X^\mu \equiv \Delta_\mu^\sigma X_{;\sigma}^\mu$) on both sides of the middle term in Eq. (7), one gets:

$$W_\nu \nabla_\mu \pi^{\mu\nu} + \pi^{\mu\nu} \nabla_{\langle\mu} W_{\nu\rangle} + 2W^\mu \nabla_\mu \Pi + 2\Pi \nabla_\mu W^\mu \equiv 0. \quad (10)$$

Following [7] let us now introduce the usual thermodynamic (positive) coefficients, $\beta_{0,1,2}$, for the, respectively, scalar, vector and tensor dissipative contributions to the entropy current (8), and the viscous/heat coupling coefficient α (all with dimension $[\text{GeV}]^{-4}$ to ensure that dimensions of identities (9) and (10) are the same as the dimension of $T[su^\mu]_{;\mu}$ in Eq. (8)). Multiplying (9b), (9c), (9a) and (10) by the, respectively, β_0 , β_1, β_2 and $\Gamma\alpha$, and combining them together one gets the following identity:

$$\begin{aligned} & \frac{2}{3} \beta_1 \Gamma W_\mu \frac{dW^\mu}{d\tau} + \alpha \Gamma W^\nu \nabla_\mu \pi_\nu^\mu + 2\alpha \Gamma W^\mu \nabla_\mu \Pi \\ & - (\beta_0 - \beta_1 - \beta_2) \Pi \frac{d\Pi}{d\tau} + 2\alpha \Gamma \Pi \nabla_\mu W^\mu \\ & - \frac{1}{6} \beta_2 \pi_{\mu\nu} \frac{d\pi^{\mu\nu}}{d\tau} + \alpha \Gamma \pi^{\mu\nu} \nabla_{\langle\mu} W_{\nu\rangle} \\ & + (\beta_0 + \beta_1) \Pi \Gamma \frac{dw}{d\tau} + \beta_0 w \Pi \frac{d\Gamma}{d\tau} = 0. \end{aligned} \quad (11)$$

The q -entropy current is connected with su^μ and W^μ/T ,

$$s_q u_q^\mu = \frac{1}{1 + \gamma} \frac{T}{T_q} \left(su^\mu + \frac{W^\mu}{T} \right). \quad (12)$$

Applying the four divergence to both sides of Eq. (12) one gets (with $\phi_\mu \equiv \partial_\mu \gamma / (1 + \gamma) + \partial_\mu T_q / T_q$ and $\tilde{\phi} \equiv u_{;\mu}^\mu - u^\mu \phi_\mu$):

$$W_{;\mu}^\mu = -\frac{dw}{d\tau} - w\tilde{\phi} + W^\mu \phi_\mu, \quad (13)$$

from which and Eq. (6) one has

$$\Gamma \frac{dw}{d\tau} = \Gamma \left(\frac{du_\mu}{d\tau} W^\mu - \nabla_\mu W^\mu \right) - \Pi \tilde{\phi} + \Gamma W^\mu \phi_\mu \quad (14)$$

(with $W_{;\mu}^\mu = \nabla_\mu W^\mu + u_\mu \frac{dW^\mu}{d\tau}$ and $u_\mu W^\mu = 0$). Adding now Eq. (11) to (8) and using Eq. (14) one obtains

$$T[su^\mu]_{;\mu} = -\Pi \Upsilon - W^\mu \Phi_\mu + \pi^{\mu\nu} \Psi_{\mu\nu} \quad (15)$$

with Υ , Φ_μ and $\Psi_{\mu\nu}$ defined as ($\beta_0^* \equiv \beta_0 - \beta_1$, $\beta_1^* \equiv \beta_1 + \beta_0$, $\alpha^* \equiv \frac{1}{\Pi\Gamma} + \beta_1^*$):

$$\begin{aligned} \Upsilon &= (\beta_0^* - \beta_2) \frac{d\Pi}{d\tau} + u_{;\mu}^\mu \\ &+ \kappa \beta_1^* \tilde{\phi} \Pi - (2\alpha - \alpha^*) \Gamma \nabla_\mu W^\mu, \end{aligned} \quad (16a)$$

$$\begin{aligned} \Phi_\mu &= -\frac{(\beta_1^* - \beta_0^*)}{3} \Gamma \frac{dW_\mu}{d\tau} - (1 + \alpha^* \Pi \Gamma) \frac{du_\mu}{d\tau} - \beta_1^* \Pi \Gamma \phi_\mu \\ &+ \frac{(\beta_1^* + \beta_0^*)}{6} \frac{d\Gamma}{d\tau} W_\mu - \alpha \Gamma \nabla_\lambda \pi_\mu^\lambda - 2\alpha \Gamma \nabla_\mu \Pi, \end{aligned} \quad (16b)$$

$$\begin{aligned} \Psi_{\mu\nu} &= -\frac{\beta_2}{6} \frac{d\pi_{\mu\nu}}{d\tau} + u_{\langle\mu;\nu\rangle} \\ &- \frac{\beta_1^*}{6} (1 - \kappa) \tilde{\phi} \pi_{\mu\nu} + \alpha \Gamma \nabla_{\langle\mu} W_{\nu\rangle}. \end{aligned} \quad (16c)$$

The arbitrary constant $\kappa \in (0, 1)$ appearing in Eqs. (16a) and (16c) is due to an ambiguity originating from the last NexDC relation in Eq. (7) which allows us to write

$\Pi^2 = \kappa\Pi^2 + (1-\kappa)\pi^{\mu\nu}\pi_{\mu\nu}/6$. To ensure that $[su^\mu]_{;\mu} \geq 0$, we assume now (following the standard 2nd order theory [7]) a linear relationship between the thermodynamic fluxes $\Pi, W^\mu, \pi^{\mu\nu}$ and the corresponding thermodynamic forces, $\Upsilon, \Phi_\mu, \Psi_{\mu\nu}$, with the usual transport coefficients ζ, λ, η , respectively. We have then

$$T[su^\mu]_{;\mu} = \frac{\Pi^2}{\zeta} - \frac{W^\mu W_\mu}{\lambda T} + \frac{\pi^{\mu\nu}\pi_{\mu\nu}}{2\eta} \quad (17)$$

and our q -hydrodynamics leads to the following constitutive equations of the corresponding d -hydrodynamics:

$$\Pi + \tau_\Pi \frac{d\Pi}{d\tau} = -\zeta u^\mu_{;\mu} - \kappa\beta_1^* \zeta \tilde{\zeta} \Pi + l_{\Pi W} \nabla_\mu W^\mu, \quad (18a)$$

$$W_\mu + \tau_W \Delta_\mu^\sigma \frac{dW_\sigma}{d\tau} = \lambda T \left[-(1 + \alpha^* \Pi \Gamma) \frac{du_\mu}{d\tau} - \beta_1^* \Pi \Gamma \phi_\mu + \frac{(\beta_0^* + \beta_1^*)}{6} \frac{d\Gamma}{d\tau} W_\mu \right] - l_{W\pi} \nabla_\sigma \pi^\sigma_\mu - l_{W\Pi} \nabla_\mu \Pi, \quad (18b)$$

$$\pi_{\mu\nu} + \tau_\pi \Delta_\mu^\rho \Delta_\nu^\sigma \frac{d\pi_{\rho\sigma}}{d\tau} = 2\eta u_{\langle\mu;\nu\rangle} - \frac{\beta_1^*}{3} (1 - \kappa) \eta \tilde{\zeta} \pi_{\mu\nu} + l_{\pi W} \nabla_{\langle\mu} W_{\nu\rangle}, \quad (18c)$$

with the corresponding relaxation times $\tau_\Pi \equiv \zeta(\beta_0^* - \beta_2)$, $\tau_W \equiv \frac{1}{3}(\beta_1^* - \beta_0^*)\Gamma\lambda T$ and $\tau_\pi \equiv \frac{1}{3}\beta_2\eta$ and with the corresponding heat-viscous coupling lengths $l_{\Pi W} = \zeta(2\alpha - \alpha^*)\Gamma$, $l_{W\Pi} = 2\alpha\Gamma\lambda T$, $l_{W\pi} = \alpha\Gamma\lambda T$ and $l_{\pi W} = 2\eta\alpha\Gamma$. To ensure the positivity of relaxation times we must have $\beta_0 > \beta_1 + \beta_2$. Actually, one can also express coefficients $\beta_{0,1,2}$ and α in terms of the corresponding relaxation times and transport coefficients:

$$\begin{aligned} \beta_0 &= \frac{\tau_\Pi}{\zeta} + \frac{3\tau_\pi}{\eta} + \frac{3}{2} \frac{\tau_W}{\Gamma\lambda T}, & \beta_1 &= \frac{3}{2} \frac{\tau_W}{\Gamma\lambda T}, \\ \beta_2 &= \frac{3\tau_\pi}{\eta}, & \alpha &= \frac{l_{W\pi}}{\Gamma\lambda T} = \frac{l_{\pi W}}{2\Gamma\eta}. \end{aligned} \quad (19)$$

To summarize this part, under the conditions mentioned above, and with the help of the dissipative tensor relations, Eq. (7) and q -entropy conservation, one can obtain the corresponding constitutive equations of d -hydrodynamics, Eq. (18). They include the relaxation times and heat-viscous coupling lengths in a quite natural way. Since the original perfect q -hydrodynamics Eq. (1) does not contain any space-time scale, it is than natural that the NexDC conjecture does not introduce any definite relaxation time or viscous-heat coupling length scale. However, as seen in Eq. (18), the d -hydrodynamics obtained from the q -hydrodynamics by using the NexDC relations, possesses the causal property of the corresponding d -hydrodynamics. Let us close noticing that this d -hydrodynamics predicts that

$$2l_{W\pi} = l_{W\Pi}, \quad \frac{l_{W\pi}}{\lambda T} = \frac{l_{\pi W}}{2\eta}; \quad (20)$$

$$\frac{1}{\Pi/s} + \frac{(l_{\Pi W} + \Gamma\tau_\Pi)}{\zeta/s} = \frac{(l_{\pi W} - 3\Gamma\tau_\pi)}{\eta/s} - \frac{\tau_W}{\lambda T/s}. \quad (21)$$

As an example, let us now apply the NexDC to a q -hydrodynamics with the Bjorken type scaling initial conditions [20]. In this case the equation of motion Eq. (1) for an q -ideal fluid has a very simple form:

$$\frac{d\varepsilon_q(\tau)}{d\tau} = -\frac{\varepsilon_q(\tau) + P_q(\tau)}{\tau}, \quad (22)$$

with four velocity field $u_q^\mu = (1, 0, 0, 0)$ (we use metric $g_{\mu\nu} = (g_{\tau\tau}, g_{xx}, g_{yy}, g_{\eta\eta}) = (1, 0, 0, -\tau^2)$, where $\tau = (t^2 - z^2)^{1/2}$ and $\eta^* = \frac{1}{2} \ln(t+z)/(t-z)$). The q -energy density and q -pressure are defined, respectively, as $\varepsilon_q = u_{q\mu} \mathcal{T}_q^{\mu\nu} u_{q\nu}$ and $P_q = -\frac{1}{3} \mathcal{T}_q^{\mu\nu} \Delta_{q\mu\nu}$, by using the q -energy momentum tensor [13], $\mathcal{T}_q^{\mu\nu} = \frac{1}{(2\pi\hbar)^3} \int \frac{d^3p}{p^0} [f_q(x, p)]^q$, with the $f_q(x, p)$ given by Tsallis distribution,

$$f_q(x, p) = \left[1 - (1 - q) \frac{p_\mu u_q^\mu}{k_B T_q(x)} \right]^{1/(1-q)}. \quad (23)$$

The corresponding EoS, $P_q = P_q(\varepsilon_q)$, is obtained by eliminating the common parameter T_q from both the $P_q(T_q)$ and $\varepsilon_q(T_q)$. Fig. 1 shows the proper time evolution of T_q, T (notice that, because of Eq. (2), they are not independent) and the bulk pressure Π for the relativistic π gas (with mass $m = 0.14$ GeV) distributed according to Eq. (23). The q -hydrodynamical evolution starts at




FIG. 1: Proper time evolution of T_q, T and Π for $q = 1.00$ – 1.20 . The q -hydrodynamical evolution starts at $\tau_0 = 1.0$ fm. The initial temperatures, $T_q(\tau_0)$ are chosen in such way as to have $T_0 = 0.2$ GeV for all values of q . The dotted line in (b) is obtained for a fluid composed of massless particles (in this case there is no q -dependence).

proper time $\tau_0 = 1.0$ fm with the q -temperature $T_q(\tau_0)$ chosen such that $T_0 \equiv T(\tau_0) = 0.2$ GeV. As seen in Fig. 1, although T_q depends on q , the $T(\tau)$ does not (it scales with q). This is interesting because in the usual d -hydrodynamics cooling behavior (i.e., T) is affected by the viscous effects. However, in our case the q -dependence of Eq. (22) is only implicit and the corresponding EoS depends only very weakly on q in the high temperature region (cf., Fig. 1 in [13]). On the other

hand, it depends on the mass of particles which composed our fluid. The bulk pressure Π decreases monotonously with τ and increases with the non-extensive parameter q .

We close this example by discussing the corresponding Reynolds number [21, 22]. Separating in Eq. (22) dissipative and non-dissipative terms by using the Reynolds number R_e [7] we get

$$\frac{d\varepsilon}{d\tau} + \frac{\varepsilon + P}{\tau} = \frac{(\varepsilon + P)}{\tau R_e}, \quad (24)$$

In q -hydrodynamics with the NexDC conjecture

$$R_e^{-1} = \frac{3\tau}{\varepsilon + P} \left(\frac{\Pi}{\tau} + \frac{d\Pi}{d\tau} \right). \quad (25)$$

Fig. 2 shows examples of proper time evolution of the inverse of Reynolds number obtained in Eq. (25). For comparison, the maximal values of R_e^{-1} quoted in [9] for $\tau_0 = 1$ fm and $T_0 = 0.3$ GeV are $R_e^{-1} \approx 0.065$ (for $\eta/s = 0.08$ and at $\tau \approx 2$ fm) and $R_e^{-1} \approx 0.21$ (for $\eta/s = 0.30$ and $\tau \approx 4$ fm). Our R_e^{-1} remains smaller than these values, nevertheless we observe a similar maximum of R_e^{-1} at finite proper time, $\tau \sim 10$ fm. Notice that one can rewrite Eq. (24) as $d\varepsilon/d\tau = (R_e^{-1} - 1)w/\tau$, which, in the case of $R_e^{-1} > 1$, may indicate the instability of our solution for large τ [7]. However, as one can see in Fig. 2, our q -hydrodynamical model with q -scaling solution gives very small values of R_e^{-1} , $R_e^{-1} \ll 1$. This, according to [22, 23], may guarantee the stability of the corresponding d -hydrodynamics obtained.




FIG. 2: Proper time evolution of R_e^{-1} obtained from q -hydrodynamics for two different initial temperatures (panels (a) and (b)) and for different values of the parameter q . Solid lines are for, respectively, $q = 1.05, 1.10$ and 1.20 , dashed lines are for $q = 1.30 - 1.50$. The same EoS has been used as in Fig. 1.

To summarize: we have demonstrated that d -hydrodynamics obtained from the perfect q -hydrodynamics (by means of the NexDC conjecture)

preserves its original causality. We believe then that q -hydrodynamics can serve as a phenomenological model of d -hydrodynamic with parameter q describing summarily non ideality of the hadronic fluid. We have also shown that the corresponding inverse of the Reynolds number, $1/R_e$, may be small enough to ensure a stable hydrodynamical evolution of the corresponding strongly interacting quark/hadronic matter.

Partial support (GW) from the Ministry of Science and Higher Education under contract 1P03B02230 is acknowledged.

* Electronic address: osada@ph.ns.musashi-tech.ac.jp

† Electronic address: wilk@fuw.edu.pl

- [1] T. Hirano, N. Kolk and A. Bilandzic, arXiv:0808.2684, to appear in Springer Lecture Notes in Physics.
- [2] T. D. Lee, Nucl. Phys. **A750**, 1 (2005).
- [3] T. Hirano and M. Gyulassy, Nucl. Phys. A **769**, 71 (2006).
- [4] C. Eckart, Phys.Rev. **58**, 919 (1940).
- [5] W. A. Hiscock and L. Lindblom, Phys. Rev. **D 31**, 725 (1985); **D 35**, 3723 (1987).
- [6] W. Israel, Ann. Phys. (N.Y.) **100**, 310 (1976); J. M. Stewart, Proc. R. Soc. London, Ser. A **357**, 59 (1977); W. Israel and J. M. Stewart, Ann Phys. (N.Y.) **118**, 341 (1979).
- [7] A. Muronga, Phys. Rev. **C 69**, 034903 (2004) and **C 76**, 014910 (2007).
- [8] U. Heinz, H. Song and A.K. Chaudhuri, Phys. Rev. **C 73**, 034904 (2006); H. Song and U. Heinz, Phys. Lett. B **658**, 279 (2008).
- [9] R. Baier and R. Romatschke and U. A. Wiedemann, Phys. Rev. **C 73**, 064903 (2006).
- [10] T. Koide, G. S. Denicol, Ph. Mota and T. Kodama, Phys. Rev. **C 75**, 034909 (2007);
- [11] A. K. Chaudhuri, Phys. Rev. **C 74**, 044904 (2006).
- [12] P.Ván and T.S.Biro, Eur. Phys.J. Special Topics **155**, 201 (2008); T.S. Biro, E. Molnar and P. Ván, Phys. Rev. **C 78**, 014909 (2008) and references therein.
- [13] T. Osada and G. Wilk, Phys. Rev. **C 77**, 044903 (2008) and Prog. Theor. Phys. Suppl. **174**, 168 (2008).
- [14] C. Tsallis, J. Stat. Phys. **52** 479 (1988). For updated bibliography on this subject see <http://tsallis.cat.cbpf.br/biblio.htm>.
- [15] A. Lavagno, Phys. Lett. **A301**, 13 (2002).
- [16] J. A. S. Lima, R. Silva and A. R. Plastino, Phys. Rev. Lett. **86**, 2983 (2001).
- [17] T. Kodama, H. T. Elze, C. E. Aguiar and T. Koide, Europhys. Lett. **70**, 439 (2005).
- [18] T.S. Biró and G. Purcsel, Phys. Rev. Lett. **95** 162302 (2005); Phys. Lett. A **372**, 1174 (2008).
- [19] T. Sherman and J. Rafelski, Lecture Notes in Physics **633** (2004) 377.
- [20] J. D. Bjorken, Phys. Rev. **D27**, 140 (1983).
- [21] G. Baym, Nucl. Phys. **A418**, 525c (1984).
- [22] H.Kouno, M. Maruyama, F. Takagi, and K. Saito, Phys. Rev. **D 41**, 2903 (1990).
- [23] G.S.Denicol, T. Kodama, T. Koide and Ph. Mota, Phys. Rev. **C 78**, 034901 (2008).